

**A Dendrochronological Analysis in Canadian Prairie Shelterbelts:  
MacMillan's Farm II**



Emma L. Davis and Colin P. Laroque

MAD Lab Report 2012-11

Mount Allison University, Department of Geography and the Environment  
Mount Allison Dendrochronology Lab

## **Table of Contents**

Abstract.....	1
Introduction.....	2
Site Information.....	2
Methods.....	3
Results.....	3
Conclusion.....	5

## **Abstract**

To determine the carbon storage capacity of shelterbelt trees and their response to climate variables, the Mount Allison Dendrochronology Lab conducted a tree-ring analysis on nine of the most commonly planted shelterbelt species in the Canadian Prairies. Traditional cross-dating and climate analyses techniques were used to reveal a variety of temporal patterns in tree-growth. At MacMillan's Farm hybrid poplar samples were collected for analyses and it was determined that the oldest trees are aged 98 years at the diameter at breast height.

## **Introduction**

In the summer of 2011, the Mount Allison Dendrochronology Lab travelled to Saskatchewan to sample shelterbelt trees as part of the Agricultural Greenhouse Gas Program (in association with the University of Saskatchewan). The objective of the larger project is to determine the carbon storage capacity of shelterbelt trees in order to determine their ability to off-set carbon emissions and act as potential carbon credits for landowners.

Samples for this project were collected around south-central Saskatchewan throughout the summer of 2011 for a dendrochronological (tree-ring) analysis in an effort to reveal the climatic factors that have had the greatest impact on annual-tree growth for the tested species. The objective of this sampling was to determine the age and growth patterns of nine of the most commonly planted shelterbelt species. As a landowner and thus a stakeholder in this project, we would like to provide you with the results of our findings on your property.

## **Site Information**

**MAD Lab Site Code:** 11ML

**Date:** MAY 28, 2011

**Site Name:** MACMILLAN FARM II

**Site Contact Info:** IAN, JUDY MACMILLAN

**Latitude:** N 52° 06' 37.4"

**Longitude:** W 107° 18' 34.8"

**UTM:** 184, 583

**UTM Zone:** 13 U

**MASL:** 539 m

**Satellites:** 5

**NAD:** 83

**Elevation:** 539 m

**Easting:** 0341843

**Northing:** 5775834

**Species Common Name:** Hybrid Poplar

**MAD Lab Species Code:** N00


## Methods

Forty tree cores were sampled from each species using a 5.1 mm increment borer. The samples were stored in plastic straws and transported to the Mount Allison Dendrochronology Lab for analysis. The samples were glued into slotted mounting boards, and then sanded and buffed to a fine polish in order to reveal the tree rings. Annual-growth rings were counted and measured using a mounted measuring stage and 60X microscope. The individual core measurements were crossdated (pattern-matched) against other cores within their group to establish the years that had increased or suppressed radial growth. A master chronology was established for each species at each site, demonstrating the overall tree-growth patterns through time.

Annual tree-ring measurements were then compared to historical temperature and precipitation data from the Saskatoon climate station in order to determine the major environmental factors influencing the tree's growth. The resulting statistical correlations allow us to infer the climate variables that play the most significant role in the growth of each shelterbelt species.


## Results

The oldest sampled hybrid poplar trees on the property were found to be 98 years old at breast height, suggesting that they were planted in the early 1910's (Fig. 1). The mean ring-width measurement was determined to be 2.14 mm.



**Fig. 1** - Master chronology for hybrid poplar (11MLN00) at MacMillan's Farm. A standardized measurement of one indicates an average year of growth, while any value above or below one indicates a year of above or below average growth.

The main climate factors controlling the majority of the year-to-year ring-width variability were previous August temperature (negative correlation), current July temperature (negative), and previous September precipitation (positive). Warm August temperatures during the previous year have a negative effect on radial tree growth of the current year due to moisture depletion from the soil. This prevents the trees from being able to store up resources for the following growing season. With similar reasoning, September precipitation of the previous year encourages growth during the current year, as increased precipitation replenishes moisture to the soil and allows the trees to take up and store nutrients for the following spring. July temperatures during the current growth season are negatively correlated to tree growth; once again, the soil is depleted of moisture, causing a decrease in radial growth.



**Fig. 2** Results of the climate analysis comparing annual tree-ring growth to historical temperature and precipitation variables from Saskatoon. Bars represent the degree of correlation between growth and the climate variable, with anything surpassing the linear thresholds being considered significantly correlated. The uppercase letters (i.e. APR T) present variables from the previous year (for example, the conditions of the previous fall often have an impact on tree growth during the current year).

## **Conclusion**

The results of these analyses have proved useful for determining the significant climatic variables influencing the annual growth of hybrid poplar in shelterbelts in south-central Saskatchewan. The data obtained through this study will aid in inferring the future growth trends of shelterbelt species under different future climate change scenarios. The eventual aim is to use this information to quantify the amount of carbon sequestered by each shelterbelt tree on an annual basis to demonstrate their potential as carbon credits.

This research was conducted at the Mount Allison Dendrochronology Lab in Sackville, New Brunswick. Any questions regarding the findings of this report should be directed to:

### **Dr. Colin Laroque**

Mount Allison Dendrochronology Lab

RoomG3, Avard-Dixon Building

Mount Allison University

144MainStreet

Sackville, NB E4L 1A7

[claroque@mta.ca](mailto:claroque@mta.ca)

506-364-2390

Thank you for your participation in this project!