

A Dendrochronological Analysis of Canadian Prairie Shelterbelts: Perry Farm



Mistik Askiwin
ᓂᓴᓴᓴ ᓴᓴᓴᓴ
Dendrochronology Lab

Scott Wood, Jacqueline Carverhill, Jason Maillet,
Nicole Marleau, Victoria Millette, and Colin P. Laroque.

MAD Lab Report 2014-10

University of Saskatchewan – Department of Soil Sciences

Mistik Askiwin Dendrochronology Lab

Table of Contents

| | |
|-----------------------------|---|
| Abstract..... | 2 |
| Introduction..... | 3 |
| Site Information..... | 3 |
| Methods..... | 3 |
| Results and Discussion..... | 4 |
| Conclusion..... | 6 |

Abstract

As a part of the Agricultural Greenhouse Gases Program, which seeks to determine the carbon sequestration capabilities of shelterbelt trees and their response to climate and climate change, the Mistik Askiwin Dendrochronology Lab conducted a tree-ring analysis on shelterbelt trees throughout Saskatchewan. Using dendrochronological cross-dating techniques and climate analysis, radial growth chronologies were established, and the relationship between tree growth and climate was examined. At the Perry's site, green ash samples were collected from twenty trees, and the oldest samples were 45-years old with an average age of 39. The main climatic factors driving tree growth at this site are previous-year July temperature, and current year May, and June precipitation.

Introduction

The Mistik Askiwin Dendrochronology Lab (MAD Lab) located at the University of Saskatchewan, is currently involved in a project for the Agricultural Greenhouse Gases Program (AGGP), which is investigating the capability of shelterbelt trees to store carbon. The carbon storage capability of these trees will inform their ability to off-set carbon emissions and potentially act as carbon credits. The objective of the larger project is to determine the current and future capacity of carbon sequestration in these shelterbelt trees.

In the summer of 2013, samples for this project were collected across most of Saskatchewan. These samples were used as a part of the larger study, looking at shelterbelt tree growth over time since the trees were planted. As a landowner, and therefore a stakeholder in this project, we would like to provide you with the results from our findings on your property.

Site Information

MAD Lab Site Code: 13LLM00

Date: June 2, 2013

Site Name: Perry Farms

Site Contact Info: Sheldon and Lillian Perry

Latitude: 51° 30' 19.2"

Longitude: -104° 10' 31.1"

UTM: 0557237 5706346

UTM Zone: 13U

MASL (m above sea level): 646m

Satellites: 7

Precision: ± 5m

Species Common Name: Green Ash

MAD Lab Species Code: M00

Methods

The MAD Lab sampled twenty green ash trees, using a 5.1 mm increment borer to take two core samples from each tree at approximately breast height. These samples were stored in plastic straws and taken back to the MAD Lab in Saskatoon, Saskatchewan for analysis. The samples were glued into slotted mounting boards and labeled with the appropriate site code. The samples were sanded with progressively finer sandpaper (60 to 600 grit) and then buffed in order to reveal the cell structure of the tree rings. The annual-growth rings were measured under a microscope using a Velmex stage system with a precision of 0.001 mm. The measurements from each core created a growth pattern, which could then be matched against the other cores from that site, in order to create a master chronology, which would demonstrate the overall tree

growth patterns through time. The master chronologies were then standardized to remove age related and biological growth trends, providing a cleaner signal.

The resulting standardized growth chronology was then used to determine the environmental factors influencing the tree's growth. Annual tree-ring measurements were compared to historical climate data from the Kelliher climate station, using the program DendroClim. The program provides statistical correlations, which allow us to identify which climate variables most significantly influence the growth of the trees at each site.

Results and Discussion

The oldest tree was found to be 45-years old at breast height, while the average age of all trees at the site was 39. This suggests they were planted in the late-60's, which is in agreement with the database provided from the Prairie Farm Rehabilitation Association, which indicates that green ash were sent there in 1967. The average raw ring-width measurement was determined to be 2.03 mm (see Figure 1 for the standardized growth of the tree over time). The climate data from the Kelliher station indicated that temperature in July and November of the previous year, and spring precipitation in the growth year (May and June) were the strongest climate variables affecting the tree growth (see Figures 2 and 3).




Figure 1: Master chronology for green ash at the Perry site. Standardized measurements of 1.00 indicate an average year of growth (in this case, associated with a raw ring-width of 2.03 mm), while any value above or below 1.00 indicate a year of growth that deviates from the average.




Figure 2: Results of the climate analysis comparing annual tree-ring growth to historical temperature variables from Kelliher, SK. The bars represent the degrees of correlation between the tree growth and the climate variable. The places where the bars cross the linear threshold are considered significantly correlated, marked by a star. The uppercase letters (ie., APR) label the previous years' variables.




Figure 3: Results of the climate analysis comparing annual tree-ring growth to historical precipitation variables from Kelliher, SK.

Conclusion

The results of this analysis help to strengthen our record of green ash growth over time within Saskatchewan's southeast. They also provide us with an understanding of the important climate variables in southeast Saskatchewan, in this case temperature in the previous-year July, and precipitation in May and June of the current year. The data used from this site will be used in future studies, which will attempt to determine future growth trends and the amount of carbon sequestered by green ash to determine its potential and viability in carbon sequestration.

This research was conducted at the MAD Lab in Saskatoon, Saskatchewan, and funded through the AGGP. Any questions regarding the findings of this report should be directed to:

Dr. Colin Laroque

Mistik Askiwin Dendrochronology Lab
Room 5C74, Agriculture Building
University of Saskatchewan
51 Campus Drive
Saskatoon, SK
S7N 5A8
Colin.Laroque@usask.ca
(306) 966-2493